

Perioperative Assessment

Stanley Kurek, DO, FACS

Associate Professor of Surgery

UTMCK Dept. of Surgery

Goals

- Understand how to estimate peri-operative CV risk
- Know when to perform stress testing preoperatively
- Learn how to reduce risk perioperatively in those at higher risk

Key Points

1. Extensive testing is rarely needed to determine risk
2. Evaluation/Testing not needed if:
 - a. Low risk surgery
 - b. Good functional capacity and no cardiac symptoms
 - c. No clinical risk factors


Key Points

4. Revascularization (surgery or PCI) should be considered only if standard indications are present
5. If PCI to be done, delay before non-cardiac surgery should be as follows:
 - POBA: 14 days
 - BMS: 30-45 days
 - DES: > 365 days

Key Points

6. Cardiac complications (both ischemia and infarction) are often manifested by:
 - a. Confusion, other MS changes
 - b. Hypotension
 - c. Dyspnea, heart failure
7. Cardiac complications tend to occur postoperatively and not intraoperatively, with a peak incidence on POD # 2-3

Key Points



ST Depression

Ref: Mangano, et al, JACC, 1991

Key Points

8. Outcomes in high risk patients optimized with:
 - a. Beta blockers
 - b. Aggressive pain control
 - c. Avoidance of severe anemia
 - d. Normothermia
 - d. Vigilant monitoring

Perioperative Risk

- Patient
 - Underlying disease
 - Physiologic Reserve
- Procedure
 - Risk Classification
- Anesthetic Risk
- Environment

Patient-associated Factors

- Underlying (comorbid) conditions
 - More comorbidity = greater risk
 - Ischemic heart disease
 - CHF
 - Diabetes
 - Renal Insufficiency
 - Low serum albumin

Patient-associated Factors (cont.)

- Undiagnosed hypothyroidism
- Hepatitis
 - Acute, 10% mortality
- Cirrhosis
- Obesity

Diabetes

- “Heroic” efforts to control BS
- Decreased postop M&M significantly in cardiac surgery
- Must be aware of hypoglycemia
- Poor wound healing
- Postop infections
- Diabetic comorbidities

Cerebral Vascular Disease

- Stroke Risk
- Hypertensive vascular disease
- Associated comorbidities

Obesity

- BMI (wt/ht)(m²)
 - BMI < 25 normal
 - BMI 25-30 overweight
 - BMI 31-40 obese
 - BMI > 40 morbidly obese

Obesity

- Decreased pulmonary reserve
 - Decreased FRC
- Wound infections
- Anesthesia difficulties:
 - Intubation
 - Venous access
 - Aspiration
- Thrombophlebitis
- Association with DM, CV disease and HTN

Patient Factors

- Renal function:
 - Approximately 1% decline in functional nephrons per year after age 40
- Pulmonary function:
 - refer to text

Overview


- Epidemiology
- Risk Assessment
- Preoperative Testing
- Postoperative Management to Reduce Risk
- Frequently asked questions
- Case studies

Epidemiology

- 43.9 million inpatient procedures annually
- CV complications are the leading cause of morbidity and mortality following surgery
 - Rates among all comers: 2%
 - >3 risk factors: 11%
- 20 Billion dollar annual cost


Source: CDC 2003 National Hospital Discharge Survey -
Published July 8, 2005

Epidemiology




Source: CDC 2003 National Hospital Discharge Survey -
Published July 8, 2005

Distribution of Procedure by Gender



Women



Men

Source: CDC 2003 National Hospital Discharge Survey - Published July 8, 2005

Triggers

- Surgical Trauma
- Anesthesia/analgesia

- Surgical Trauma
- Anesthesia/analgesia

- Surgical Trauma
- Anesthesia/analgesia
- Intubation/extubation
- Pain
- Hypothermia
- Bleeding/anemia
- Fasting

- Anesthesia/analgesia
- Hypothermia
- Bleeding/anemia

Inflammatory State

↑TNF- α
↑IL-1
↑IL-6
↑CRP

Plaque fissuring

Hypercoagulable State

↑ PAI-1
↑ Factor VII
↑ Platelet reactivity
↓ antithrombin III

Stress State

↑ catecholamine and cortisol levels

Coronary artery shear stress

Plaque fissuring

Hypoxic State

↓ oxygen delivery

↑ BP
↑ HR
↑ FFAs
↑ relative insulin deficiency

↑ Oxygen demand

Acute Coronary Thrombus

Myocardial Ischemia

Perioperative Myocardial Infarction

Men and woman are not the same



Gender Differences in Heart Disease

- Woman get it at a later age
- Woman are less likely to manifest with “typical” symptoms
- Women have worse outcomes in cardiac intervention
- Women (most) don't have wives to take care of them!

Overview

- Epidemiology
- **Risk Assessment**
- Preoperative Testing
- Postoperative Management to Reduce Risk
- Frequently Asked Questions
- Case Studies

Approaches to Risk Assessment

1. ASA/Dripps
 2. Goldman Multifactorial Index
 3. Detsky Modified Index
 4. Revised Risk Index
 5. ACC/AHA Task Force Recommendations
- Quantitative
- Strategic
-
- The diagram consists of a list of five risk assessment approaches on the left. A large white curly bracket on the right groups the first four items (ASA/Dripps, Goldman Multifactorial Index, Detsky Modified Index, and Revised Risk Index) under the label 'Quantitative'. A smaller white curly bracket on the right groups the fifth item (ACC/AHA Task Force Recommendations) under the label 'Strategic'. The labels 'Quantitative' and 'Strategic' are written in yellow text.

Dripps/ASA Classification

Class	Systemic Disturbance	Mortality*
1	Healthy patient with no disease outside of the surgical process	<0.03%
2	Mild-to-moderate systemic disease caused by the surgical condition or by other pathologic processes	0.2%
3	Severe disease process which limits activity but is not incapacitating	1.2%
4	Severe incapacitating disease process that is a constant threat to life	8%
5	Moribund patient not expected to survive 24 hours with or without an operation	34%
E	Suffix to indicate an emergency surgery for any class	Increased

Goldman Risk Index

Third heart sound (S ₃)	11
Elevated jugulovenous pressure	11
Myocardial infarction in past 6 months	10
EKG: premature arterial contractions or any rhythm other than sinus	7
EKG shows >5 premature ventricular contractions per minute	7
Age >70 yrs	5
Emergency procedure	4
Intra- Thoracic, intra-abdominal, or aortic surgery	3
Poor general status, metabolic or bedridden	3

<u>Class</u>	<u>Point Total</u>	<u>None/ Minor Complication</u>	<u>Life- Threatening Complication</u>	<u>Cardiac Death</u>
Class I	0-5	99%	0.7%	0.2%
Class II	6-12	93%	5%	2%
Class III	13-25	86%	11%	2%
Class IV	≥26	22%	22%	56%

Ref: Goldman M, Caldera D, Southwick, et al: Multifactorial index of cardiac risk in non-cardiac surgical procedures. *N Engl J Med* 148:2120-2127, 1988.

ACC/AHA Guidelines

ACC/AHA 2007 Guidelines on Perioperative Cardiovascular Evaluation and Care for Noncardiac Surgery


A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines on Perioperative Cardiovascular Evaluation for Noncardiac Surgery)


Developed in Collaboration With the American Society of Echocardiography, American Society of Nuclear Cardiology, Heart Rhythm Society, Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, and Society for Vascular Surgery

J Am Coll Cardiol, 2007; 50:1707-1732

Stepwise Approach to the Pre- operative Evaluation

Stepwise Approach to Preoperative Cardiac Assessment





*Clinical risk factors = known ischemic heart disease, compensated or prior HF, diabetes, renal insufficiency, cerebrovascular disease


Importance of Surgical Urgency

Elective Surgery: Carried out at a time to suit the patient and surgeon

Urgent Surgery: Carried out within 24-hrs of admission

Emergency Surgery: Carried out within 2-hrs of admission or in conjunction with resuscitation

Non-Cardiac Surgery



Source: Evaluation of National Confidential Enquiry into Perioperative Deaths (NCEPOD)

Surgical Urgency?

Key Point: Patients undergoing urgent or emergent surgery are at higher risk of postoperative complications and require closer surveillance postoperatively.

Functional Capacity

1. Correlates with maximum oxygen uptake on treadmill testing
2. Demonstrated predictor of future cardiac events
2. Poor functional capacity may hide low threshold cardiac symptoms



Duke Activity Status Index

1 MET

Can you take care of yourself?
Eat, dress, or use the toilet?
Walk indoors around the house?
Walk a block or two on level ground at 2-3 mph or 3.2-4.8 km/h?



4 METs

Do light work around the house like dusting or washing clothes?

MET = metabolic equivalent

4 METs

Climb a flight of stairs or walk up a hill?
Walk on level ground at 4 mph or 6.4 km/h?
Run a short distance?
Do heavy work around the house like scrubbing floors or lifting or moving heavy objects?
Participate in moderate recreational activities like golf, bowling, dancing, doubles tennis, or throwing a baseball or football?




10 METs

Participate in strenuous sports like swimming, singles tennis, football, baseball, or skiing?

Assessing Risk

Active Cardiac Conditions



High Risk:

- Acute or recent MI (7-30 d)
- Unstable coronary syndrome
- Decompensated CHF
- Significant Arrhythmias
- Severe Valvular Disease

↓
~~Surgery~~

Clinical Risk Factors



Proceed Cautiously with:

- History of heart disease
- Compensated or prior CHF
- Cerebrovascular disease
- Diabetes Mellitus
- Renal Insufficiency



3 or more risk factors
& Vascular surgery

1 – 2 risk factors

Consider testing


Proceed with surgery
or consider testing

Low Risk Situations



Low Risk:

- Low risk surgery
- Good functional capacity
- No cardiac symptoms
- No “active cardiac conditions”
- No clinical risk factors



Reasonable to proceed with surgery

Surgery Related Risk

High Risk (Risk > 5%):

Emergent major operations

Aortic and other major
vascular

Peripheral vascular

Anticipated prolonged or
associated with large
fluid shifts and/or
blood loss

Intermediate Risk

(Risk < 5%):

Carotid
endarterectomy

Endovascular AAA
repair

Head and neck

Intraperitoneal and
intrathoracic

Orthopedic

Prostate

Low Risk Surgery

(Risk < 1%):


- ✓ Endoscopic procedures
- ✓ Superficial procedure
- ✓ Cataract surgery
- ✓ Breast surgery

Overview

- Epidemiology
- Risk Assessment
- Preoperative Testing
- Postoperative Management to Reduce Risk
- Frequently Asked Questions
- Case Studies

Preoperative Testing

Negative Predictive Value




Eagle et al. JACC 1996;27:910.

Preoperative Testing

Positive Predictive Value

MI or Death



ACC/AHA Recommendations

- Echocardiography:
 - Dyspnea of unknown origin (Class IIa)
 - Current or hx of HF and no echo in 12 months (Class IIa)
- 12 Lead ECG
 - Vascular surgery and 1 CRF (class I)
 - CRFs and intermediate risk surgery (class I)
 - All vascular surgery (class IIa)

ACC/AHA Recommendations

- Treadmill stress testing
 - High cardiac risk conditions
 - 3 CRFs, poor functional capacity & vascular surgery (class IIa)
- Nuclear stress testing

Which test to choose?

Most ambulatory
patients



Treadmill Stress Test

Abnormal resting
ECG (dig, LVH)



Exercise
echo or sestamibi

LBBB
Unable to exercise



DSE
Adenosine sestamibi
dipyridamole sestamibi


Overview

- Epidemiology
- Risk Assessment
- Preoperative Testing
- **Postoperative Management to Reduce Risk**
- Frequently Asked Questions
- Case Studies


Perioperative Management

- Revascularization
- Beta blockers
- Statins
- Alpha-2 agonists
- Calcium channel blockers


PCI before anticipated surgery



Timing of Surgery After PCI



Postoperative Mortality Reduction Beta-Blockers




- 200 pts undergoing non-cardiac surgery
- Random assignment to:
 - IV followed by oral atenolol or
 - Placebo
- Double-blind follow-up over 2 years

Mangano, et al. NEMJ 1996;335:1713.

Postoperative Cardiac Events In High Risk Patients

Beta-Blockade



- 173 patients undergoing vascular surgery with positive DSE

- Randomized to BB 1 week pre-op or placebo

- Followed for 30 days

Poldermans et al. NEJM 1999;341:1789.

Perioperative Beta Blockers

AHA/ACC Recommendations: 2006 Update

- Beta blockers required in recent past to control symptoms of angina or patients with symptomatic arrhythmias or hypertension
- Patients at high cardiac risk owing to the finding of ischemia on preoperative testing who are undergoing vascular surgery
- Patients undergoing vascular surgery and with identified CAD
- Vascular surgery and multiple cardiac risk factors
- Moderate or high risk surgery and multiple cardiac risk factors

Key Point: if known or suspected CAD and undergoing moderate or high risk surgery, use a beta blocker!

Perioperative Nitrates?

35

ACC/AHA 2007 Recommendations: Nitrates

Class I: None

Class IIa: None

Class IIb: Can consider in context of anesthetic plan and patient hemodynamics

0

Preop

Induction


Incision

Emerg.

PostOp


Perioperative Statins?

- 100 patients pre-op before vascular surgery
- Random assignment:
 - Atorvastatin 20 mg
 - Placebo
- Started 30 days preoperatively
- Follow-up 6 month
- Endpoint:
 - Cardiac death
 - Non-fatal MI
 - USA
 - Stroke



Characteristic	Atorvastatin (n = 50)		Placebo (n = 50)		P
	n	%	n	%	
Death from cardiac causes	1	2.0	2	4.0	1.000
Nonfatal acute myocardial infarction	3	6.0	8	16.0	.199
Unstable angina	—		1	2.0	1.000
Ischemic stroke	—		2	4.0	.495
Combined end point	4	8.0	13	26.0	.031

Perioperative Statins




Hindler, et al. Anesthesiology 2006;105:1260-72

Treatment of Anemia?

- There is a direct relationship between preoperative anemia and risk of complications (CV complications, infection, mortality), especially if known CAD
- Decline in Hbg associated with increase in mortality, especially in those with CV disease
- Benefits of transfusion have not been proven

Preoperative Hgb and Mortality


Study of Untreated Anemia



Carson, et al. Lancet. 1996;348:1055-60

Perioperative Hypothermia

■ Normothermia ■ Hypothermia



- 300 pts undergoing

ACC/AHA 2007 Recommendations: Normothermia

Class I: For most procedures except during periods in which mild hypothermia is intended for organ protection

y
double
ment
or

Key Point:

Avoid Sympathetic Stimulation in those at Risk!

- Beta blocker if able
- Limit hypothermia
- Aggressive post-operative pain control
- Avoid significant anemia

Overview

- Epidemiology
- Risk Assessment
- Preoperative Testing
- Postoperative Management to Reduce Risk
- Frequently Asked Questions
- Case Studies

FAQs

How should I handle the patient with a pacemaker or implantable defibrillator (ICD)?

- ICDs should be turned off immediate preoperative and turned on immediately postoperatively.
- Pacemakers can be left on perioperatively but should be interrogated pre- and post-op.

FAQs

How long should surgery be delayed after a myocardial infarction?

- In general, at least one month.
- In patients treated with PCI, delay based on type of treatment:
 - POBA: 14 days
 - BMS: 30 days
 - DES: 365 days

FAQs

How should I handle the patient with atrial fibrillation?

- If rate controlled, it is not a reason to delay surgery or expect problems.
- If on warfarin, should communicate with PCP or cardiologist about safety of discontinuing.

FAQs

Can anticoagulation be stopped in the patient with a mechanical heart valve?

- Low risk patients (bileaflet Aortic valve, no risk factors)
 - Stop warfarin 48-72 pre-op
 - Resume 24 hrs post-op
- High risk patients (mitral valve, aortic valve + any risk factor)
 - Bridge with UFH, starting when INR < 2

Risk factors: AF, previous thromboembolism, LV dysfunction, hypercoagulable state, older generation valve, mechanical tricuspid valve, more than one mechanical valve

Questions?